Insights from a Long-term Study of Persistent *Exxon Valdez* Oil on Katmai and Kenai Fjords National Park Shores




Gail V. Irvine

USGS Alaska Science Center

Daniel F. Mann

Univ of Alaska, Fairbanks

Jeffrey W. Short
JWS Consulting





Exxon Valdez Spill History


- March 24,1989
- 10.8 million gallons of North Slope crude oil
- Spill point: Bligh Reef in NE Prince
 William Sound






Geographical Extent of Exxon Valdez Oil Spill and Study Site Locations

 Gulf of Alaska sites






Study Sites








- McArthur Pass
- Cape Douglas
- Kiukpalik Island

- Ninagiak Island
- Cape Gull
- Kashvik



Boulder Armors and Oil Persistence





Oil Mousse





Objectives

- Monitor the persistence and degradation of oil at Gulf of Alaska beaches
- Investigate the mechanisms that allow its persistence
- 2011- Test whether oil is being released from the sites



Study Plan Components

- Quantify surface oiling (quadrats)
- Subsurface oiling (dipstones)
- Oil chemistry (GC/MS)
- Boulder stability (re-survey marked boulders)
- Deploy plastic lipophilic films ('pucks')




Surface Oil






Change in Surface Oil Cover





Dip Stones





'A' = reinserted dip stone; 'B'= dislodged dip stone Cape Douglas, 2005



Boulder Armor






Boulder Armor



Boulder Movement


Cape Douglas


- Movement of bolts set in boulders
- Circles = 2005 data
- Squares = 1999 data
- Both are relative to 1994 positions






Boulder Armors Differ







Site or Quadrat Infilling





Ninagiak




Pertinent Oil Chemistry

- Polycyclic aromatic hydrocarbons -TPAHs
- Normal alkanes



Weathering






GOA and Prince William Sound Oil Weathering

 Gulf of Alaska Beaches


TPAH

n-alkanes





Leaking Oil?







2011 Images









Results

- 16- 22 years persistence of Exxon Valdez oil at sites distant from spill origin
- As of 2005, surface oiling has declined significantly, but subsurface oiling has changed little
- Stability of boulder armors important in the continuing persistence of the oil
- As of 2005, oil has not weathered chemically, and remained compositionally similar to 11-day-old *Exxon* Valdez oil, at most sites
- Mousse formation allowed long-distance transport and enhanced preservation of only slightly weathered oil



Acknowledgements

- Funding and support from the Exxon Valdez Oil Spill Trustee Council, the U.S. Geological Survey and the National Park Service
- Thanks to: Carl Schoch, Joel Cusick, Buddy Goatcher, Jennifer Mondragon, Jeff Mondragon, Erica Madison, Yvette Gillies, Sharon Kim, Marie Larsen, Jeep Rice, Mark Carls, Larry Holland, Carson Baughman, Carissa Turner, Tom Scheer, and captains and crews of the R/V Waters, Kittiwake II, and the Puk-uk





Infilling of Quadrats









Kiukpalik: 1994 2005




Boulder Armor









Exposure of Beaches

Monitoring



Objectives

- Monitor the persistence and degradation of oil at Gulf of Alaska beaches
- Investigate the mechanisms that allow its persistence
- 2011- Test whether oil is being released from the sites



